

Porous polymer monolithic columns with gold nanoparticles as an intermediate ligand for the separation of proteins in reverse phase-ion exchange mixed mode

Terborg, Lydia

2014

Terborg , L , Masini , J C , Lin , M , Lipponen , K H , Riekkola , M-L & Svec , F 2014 , ' Porous polymer monolithic columns with gold nanoparticles as an intermediate ligand for the separation of proteins in reverse phase-ion exchange mixed mode ' , Journal of Advanced Research , vol. 6 , no. 3 , pp. 441-448 . <https://doi.org/10.1016/j.jare.2014.10.004>

<http://hdl.handle.net/10138/224240>

<https://doi.org/10.1016/j.jare.2014.10.004>

cc_by_nc_nd

publishedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

ORIGINAL ARTICLE

Porous polymer monolithic columns with gold nanoparticles as an intermediate ligand for the separation of proteins in reverse phase-ion exchange mixed mode

CrossMark

Lydia Terborg^a, Jorge C. Masini^b, Michelle Lin^c, Katriina Lipponen^d,
Marja-Liisa Riekollä^d, Frantisek Svec^{a,*}

^a The Molecular Foundry, E.O. Lawrence Berkeley National Laboratory, Berkeley, CA 94720, United States

^b Institute of Chemistry, Department of Fundamental Chemistry, University of São Paulo, C.P. 26077, 05513-970 São Paulo, Brazil

^c Department of Chemistry, University of California, Berkeley, CA 94720, United States

^d Laboratory of Analytical Chemistry, Department of Chemistry, P.O. Box 55, FIN-00014, University of Helsinki, Helsinki, Finland

ARTICLE INFO

Article history:

Received 28 September 2014

Received in revised form 15 October 2014

Accepted 16 October 2014

Available online 4 November 2014

Keywords:

Gold nanoparticles

Mixed mode

Monolith

Proteins

Separation

ABSTRACT

A new approach has been developed for the preparation of mixed-mode stationary phases to separate proteins. The pore surface of monolithic poly(glycidyl methacrylate-co-ethylene dimethacrylate) capillary columns was functionalized with thiols and coated with gold nanoparticles. The final mixed mode surface chemistry was formed by attaching, in a single step, alkanethiols, mercaptoalkanoic acids, and their mixtures on the free surface of attached gold nanoparticles. Use of these mixtures allowed fine tuning of the hydrophobic/hydrophilic balance. The amount of attached gold nanoparticles according to thermal gravimetric analysis was 44.8 wt.%. This value together with results of frontal elution enabled calculation of surface coverage with the alkanethiol and mercaptoalkanoic acid ligands. Interestingly, alkanethiols coverage in a range of 4.46...4.51 molecules/nm² significantly exceeded that of mercaptoalkanoic acids with 2.39...2.45 molecules/nm². The mixed mode character of these monolithic stationary phases was for the first time demonstrated in the separations of proteins that could be achieved in the same column using gradient elution conditions typical of reverse phase (using gradient of acetonitrile in water) and ion exchange chromatographic modes (applying gradient of salt in water), respectively.

^a 2014 Production and hosting by Elsevier B.V. on behalf of Cairo University.

* Corresponding author. Tel./fax: +1 510 486 7964.

E-mail address: fsvec@lbl.gov (F. Svec).

Peer review under responsibility of Cairo University.

Introduction

Mixed-mode chromatography refers to chromatographic methods that utilize more than one type of interaction between the stationary phase and analytes in order to achieve their separation [1,2]. While early chromatographic methods preferred stationary phases with a strictly singular functionality,

